

## "Impact of Photocatalytic Processes on Herbicide Removal from Water"

Jagdish Ram, Research Scholar, Dept. of Chemistry, Shri Khushal Das University, Pilibanga, Hanumangarh  
Prof. (Dr.) Shyamvir Singh, Dept. of Chemistry, Shri Khushal Das University, Pilibanga, Hanumangarh

### Abstract

The presence of herbicides in water sources is a major environmental concern, as they pose risks to aquatic ecosystems and human health. Photocatalytic processes, which utilize light-activated catalysts to degrade organic contaminants, have emerged as an effective solution for removing herbicides from contaminated water. This paper reviews the impact of photocatalytic processes, particularly using materials like titanium dioxide (TiO<sub>2</sub>) and zinc oxide (ZnO), on herbicide degradation. It examines key factors such as light intensity, pH, catalyst concentration, and temperature that influence degradation efficiency. The study also compares different photocatalytic materials and reaction conditions, emphasizing the practical application of this technology for large-scale water treatment.

### Introduction:

Herbicides are commonly used in agriculture to control weed growth, but their widespread application has led to significant contamination of water bodies through runoff and leaching. Conventional water treatment methods, such as filtration and chemical treatments, are often inadequate in removing herbicides due to their persistence and chemical stability. Photocatalysis, a process where light energy is used to activate a catalyst that accelerates the breakdown of pollutants, has been shown to effectively degrade organic contaminants, including herbicides, in water. Titanium dioxide (TiO<sub>2</sub>) is the most studied photocatalyst due to its high efficiency, stability, and low toxicity. However, other materials such as zinc oxide (ZnO) and composite catalysts have also gained attention for their potential in improving photocatalytic herbicide removal. This paper reviews the mechanisms, advantages, and challenges of photocatalytic herbicide degradation, providing insights into how this process can be optimized for real-world applications in water treatment.

### Literature Review:

**Park and Choi (2023)** investigate the synergistic effects of combining titanium dioxide (TiO<sub>2</sub>) with carbon-based composites for the photocatalytic degradation of herbicides. The authors discuss how carbon materials, such as graphene oxide and carbon nanotubes, enhance the photocatalytic activity of TiO<sub>2</sub> by improving electron transfer, increasing surface area, and expanding light absorption into the visible spectrum. The review emphasizes the effectiveness of these composite photocatalysts in breaking down herbicides in water, showing superior degradation performance compared to traditional TiO<sub>2</sub> alone. The study also highlights the potential for these advanced composites to offer more sustainable and efficient solutions for water purification and environmental protection.

**Gupta and Rani (2015)** compare the effectiveness of titanium dioxide (TiO<sub>2</sub>) and zinc oxide (ZnO) photocatalysts for the degradation of atrazine, a widely used herbicide, in aqueous solutions. The authors assess various parameters, such as catalyst concentration, pH, and light intensity, to evaluate the photocatalytic efficiency of both materials. The study finds that TiO<sub>2</sub> exhibited higher degradation rates compared to ZnO, primarily due to its greater stability and better photocatalytic performance under UV light. However, ZnO also showed significant degradation efficiency, especially under certain experimental conditions. This comparative study provides valuable insights into the selection of photocatalysts for effective herbicide removal from contaminated water.

### Objectives:

1. To investigate the effectiveness of photocatalytic processes in removing herbicides from water.

2. To identify key factors that influence the efficiency of photocatalytic herbicide degradation, including light source, catalyst type, pH, and temperature.
3. To compare the performance of different photocatalysts, such as TiO<sub>2</sub>, ZnO, and composite materials, in herbicide degradation.
4. To assess the impact of environmental factors, such as natural water matrices, on the photocatalytic process.
5. To evaluate the scalability and practicality of photocatalysis as a method for large-scale water treatment.

### **Methodology:**

**Selection of Photocatalysts:** Titanium dioxide (TiO<sub>2</sub>), zinc oxide (ZnO), and composite photocatalysts are chosen for their proven efficiency in the photocatalytic degradation of herbicides. TiO<sub>2</sub> is widely recognized for its high photocatalytic activity under UV light, while ZnO offers good performance under both UV and visible light. Composite materials, combining TiO<sub>2</sub> or ZnO with other substances like graphene or carbon-based materials, enhance light absorption and reduce electron-hole recombination, making them promising candidates for improved herbicide removal

### **Experimental Setup:**

A batch reactor system is utilized for the photocatalytic experiments. The reactor is designed with both UV and visible light sources, allowing for the simulation of various lighting conditions to evaluate the efficiency of herbicide degradation under different light environments. This setup ensures controlled experimentation, where the impact of light type, catalyst, and other factors on the degradation process can be accurately assessed. Various parameters such as catalyst concentration, pH, temperature, and herbicide concentration are controlled to study their effects on degradation.

### **Herbicides Used:**

The herbicides selected for the experiments are commonly found in contaminated water bodies, including atrazine, glyphosate, and 2,4-D. These herbicides are widely used in agricultural practices and pose significant environmental concerns due to their persistence in water. Their degradation through photocatalysis is critical to assessing the effectiveness of the photocatalytic process for removing harmful contaminants from aquatic environments.

### **Monitoring of Degradation:**

The degradation of herbicides is monitored using High-Performance Liquid Chromatography (HPLC) and UV-Vis spectroscopy to track the herbicide concentrations at various time intervals. These techniques provide accurate measurements of the herbicide levels, allowing for the assessment of the degradation process. Additionally, the degradation rate constant is calculated using kinetic models, which quantifies the speed at which the herbicides are removed from the water, helping to evaluate the effectiveness of the photocatalytic process.


### **Data Collection:**

Experimental data is collected on the degradation efficiency under different experimental conditions. Variables such as light intensity, pH, and temperature are varied to determine their impact on the photocatalytic process. Experimental data is gathered to assess the degradation efficiency of herbicides under varying conditions. Key variables such as light intensity, pH, and temperature are systematically altered to observe their effects on the photocatalytic process. This data helps in understanding how each factor influences the degradation rate, allowing for the optimization of experimental conditions for maximum herbicide removal efficiency.

### **Data Analysis:**

The experimental data is analyzed to assess the degradation efficiency of different herbicides under various photocatalytic conditions. First-order kinetics is typically applied to model the degradation process, with degradation rate constants calculated for each herbicide. The results

reveal the influence of key factors such as light source (UV vs. visible light), catalyst concentration, pH, and temperature on the efficiency of herbicide removal. The degradation efficiencies of TiO<sub>2</sub>, ZnO, and composite materials are compared, showing the advantages and limitations of each. Statistical analysis is conducted to evaluate the significance of each parameter in the degradation process, and optimal conditions for herbicide removal are determined.



**Figure: Removal of pesticides from water and wastewater by solar-driven photocatalysis - ScienceDirect**

### Conclusion:

Photocatalytic degradation has proven to be an effective method for removing herbicides from contaminated water. Titanium dioxide (TiO<sub>2</sub>) and zinc oxide (ZnO) are the most widely studied photocatalysts, with TiO<sub>2</sub> demonstrating high efficiency and stability under UV light. However, factors such as light intensity, catalyst concentration, pH, and temperature significantly influence the degradation rate. Composite photocatalysts have shown promising results in improving the efficiency and extending the light absorption range. Environmental factors, such as the natural water matrix (e.g., turbidity, dissolved organic matter), can affect the photocatalytic efficiency, making it essential to optimize reaction conditions for real-world applications. While photocatalysis offers a sustainable solution for herbicide removal, challenges such as catalyst stability and energy efficiency remain. Further research is needed to enhance catalyst performance and develop cost-effective, large-scale water treatment systems based on photocatalytic processes.

### References

1. Chen, D., & Zhang, Y. (2024). Photocatalytic degradation of 2,4-D herbicide in aquatic environments: A review on catalysts and mechanisms. *Environmental Science & Technology*, 58(2), 101-113. <https://doi.org/10.1021/acs.est.3c06102>
2. Park, S., & Choi, W. (2023). Photocatalytic herbicide degradation: Synergistic effects of TiO<sub>2</sub> and carbon-based composites. *Environmental Pollution*, 298, 118779. <https://doi.org/10.1016/j.envpol.2022.118779>
3. Vassilaki, A., & Makropoulos, C. (2022). Advanced photocatalytic processes for herbicide removal: A comparative study of TiO<sub>2</sub>-based photocatalysts. *Journal of Hazardous Materials*, 423, 127068. <https://doi.org/10.1016/j.jhazmat.2021.127068>
4. Jiang, H., & Li, X. (2021). Degradation of glyphosate and atrazine using ZnO-based photocatalysts under visible light. *Journal of Environmental Sciences*, 106, 93-101. <https://doi.org/10.1016/j.jes.2021.03.016>
5. Zhai, L., & Li, Z. (2020). Photocatalytic degradation of herbicides: Recent advances in photocatalytic materials and degradation mechanisms. *Environmental Chemistry Letters*, 18(1), 85-102. <https://doi.org/10.1007/s10311-019-00931-4>

6. Cheng, X., & Xu, X. (2019). Photocatalytic degradation of atrazine in water using TiO<sub>2</sub> and graphene oxide composites: A detailed study on reaction kinetics and intermediates. *Journal of Photochemistry and Photobiology A: Chemistry*, 381, 132013. <https://doi.org/10.1016/j.jphotochem.2019.132013>
7. Yang, F., & Lin, J. (2018). Photocatalytic degradation of herbicides: Role of TiO<sub>2</sub>-based materials and their photocatalytic mechanisms. *Chemical Engineering Journal*, 345, 560-570. <https://doi.org/10.1016/j.cej.2018.03.023>
8. Shakeri, M., & Zeynizadeh, B. (2017). Effect of light intensity and pH on the photocatalytic degradation of herbicides in aqueous solutions. *Journal of Environmental Chemical Engineering*, 5(6), 5677-5684. <https://doi.org/10.1016/j.jece.2017.08.029>
9. Sriram, P., & Ramaswamy, V. (2016). Photocatalytic degradation of herbicides in contaminated water: The role of doped TiO<sub>2</sub> photocatalysts. *Environmental Toxicology and Chemistry*, 35(9), 2254-2261. <https://doi.org/10.1002/etc.3475>
10. Gupta, N., & Rani, A. (2015). Photocatalytic degradation of atrazine using TiO<sub>2</sub> and ZnO photocatalysts: A comparative study. *Science of the Total Environment*, 518, 392-399. <https://doi.org/10.1016/j.scitotenv.2015.02.041>
11. Kaur, S., & Sharma, S. (2014). Removal of herbicides from water by photocatalysis using titanium dioxide. *Journal of Hazardous Materials*, 280, 45-53. <https://doi.org/10.1016/j.jhazmat.2014.07.010>
12. Xu, Y., & Zhang, W. (2013). Photocatalytic degradation of herbicides using TiO<sub>2</sub> photocatalysts: A review of the mechanisms and applications. *Journal of Environmental Sciences*, 25(8), 1562-1574. [https://doi.org/10.1016/S1001-0742\(12\)60185-5](https://doi.org/10.1016/S1001-0742(12)60185-5)
13. Wu, Z., & Zhang, X. (2012). Photocatalytic degradation of 2,4-D in aqueous solution using TiO<sub>2</sub>/graphene oxide composites under UV light. *Environmental Science & Technology*, 46(21), 11684-11691. <https://doi.org/10.1021/es302553p>
14. Shah, K., & Agarwal, M. (2011). Photocatalytic degradation of glyphosate: Application of TiO<sub>2</sub> and ZnO photocatalysts under UV and visible light. *Environmental Pollution*, 159(10), 2702-2709. <https://doi.org/10.1016/j.envpol.2011.05.031>
15. Zhang, X., & Li, Q. (2010). The photocatalytic degradation of herbicides in water using TiO<sub>2</sub> and ZnO photocatalysts. *Water Research*, 44(19), 5479-5487. <https://doi.org/10.1016/j.watres.2010.06.039>